

Geometry of unitary orbits and correlations

Kavan Modi*

Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543

(Dated: February 4, 2009)

I examine bipartite and multipartite correlations within the construct of unitary orbits. I show that the set of product states is a very small subset of set of all possible states. I further show that all bipartite states are connected to *classically correlated* states via unitary transformations. Finally I look at how entanglement looks in this picture for the two-qubit case.

PACS numbers: 02.10.Ud, 02.40.Ft, 03.65.Ud

The geometry of quantum states is an old and a rich topic [1]. In this paper, I explore the space of unitary orbits, which I define following a recent paper by Boya and Dixit [2] as the following set

$$\varrho \equiv \{U\rho U^\dagger\} \forall U, \quad (1)$$

where ρ is a density matrix and U is an unitary operator in the same space, and ϱ represents the set of all states that are connected by all possible unitary operations; hence I call it *an unitary orbit*.

An unitary transformation is a transformation of basis, which leaves the spectrum unchanged. Any two density matrices that share a spectrum, belong to a single unitary orbit; conversely if $\rho_1 \in \varrho$ and $\rho_2 \in \varrho$, then $\text{Spec}(\rho_1) = \text{Spec}(\rho_2)$. Therefore an unitary orbit is uniquely identified by the spectrum of the states it contains. Consequently, all states on an unitary orbit will also have the same scalar measures such as the von Neumann entropy and purity. On the other hand, the analysis here has more depth than a study of scalar measures alone because *two different* unitary orbits can share a single value for *entropy* or *purity* [2]. It may be tempting to simply study the space of the eigenvalues. In fact that is all I am doing, but I am taking advantage of the nice properties of unitary transformations to connect various density states.

Let me now briefly discuss the familiar example of the space of one qubit represented by the points inside of the Bloch sphere, shown on the left in Fig 1. The outer most shell of the Bloch sphere represents the set pure states, the center point represents the fully mixed state, and everything else is represented by the space inside of the sphere. The states on each of the spherical shells of the Bloch sphere are connected to each other via unitary transformations. To construct the geometry of unitary orbits, I contract each shell of the Bloch sphere into a single point. That is, all of the states that are unitarily connected are now represented by an *unitary orbit*. The geometry for the unitary orbit, for a qubit, simplifies to line, shown on the right in Fig. 1. The two end points of the line are the fully mixed state labeled as O and the set of all pure states labeled as P .

One can carry out such a procedure any dimensional systems. For a qutrit, a three-level system, the geometry




Figure 1: The states of a qubit are represented by the points inside the Bloch sphere (left). The set of states on each spherical shell are connected by a unitary transformation, therefore the sphere goes into a line (right) when the set of all qubit states are mapped to set of unitary orbits of a qubit. P represents the set of all pure states and O represents the fully mixed state.

of states is given by an eight dimensional manifold. The geometry of unitary orbit, however, is given by a triangle [2]. In general the manifold that represents the set of state for a d -dimensional system grows as $d^2 - 1$ [3], while the manifold of unitary orbit grows linearly $d - 1$. Furthermore the manifold for unitary orbits is a simplex with d vertices. One can think of each unitary orbit as a $d + 1$ dimensional manifold contracted to a point.

In this paper I want to analyze the space of unitary orbits for composite systems with three different cases in mind. First, I want look at the unitary orbits of a composite system that contain product states, followed by classically correlated states, and finally entanglement within unitary orbits. Before I proceed, I should remark that generally in quantum information theory one is interested in local operations. The unitary orbits of composite system that I consider here are due to the actions of global unitary transformations.

Theorem. The subspace of unitary orbits containing product states a $\sum_i d_i - n$ dimensional surface in a $\prod_i d_i - 1$ dimensional manifold, where d_i is the dimensions of i th subsystem.

Proof. Suppose that an unitary orbit, ϱ_p , contains a n -partite product state written as $\rho_p = \bigotimes_i^n \rho^{(i)} \equiv$

$\rho^{(1)} \otimes \rho^{(2)} \otimes \rho^{(3)} \dots \otimes \rho^{(n)}$. Diagonalize each of the subparts above by local unitary transformations, $u^{(i)}$, to yield $\rho_{pd} = \bigotimes_i^n u^{(i)} \rho_d^{(i)} u^{(i)\dagger} = \rho_d^{(i)}$. The spectrum of ρ_{pd} is simply the outer product of the spectrums of each of the subsystem. Local unitary transformations are a subset of global unitary transformations, therefore ρ_p and ρ_{pd} belong to the same unitary orbit and share the same spectrum. In fact, any state $\rho \in \mathcal{Q}_p$ has the same spectrum. Conversely, any unitary orbit that contains a product states has a spectrum that is factorable in terms of the spectrum of the subparts. Each of the subparts above has $d_i - 1$ independent parameters (eigenvalues); the total number of independent parameters for $\mathcal{Q}_p$ is simply the sum of independent parameters of each of the subpart, $\sum_i^n (d_i - 1) = \sum_i^n d_i - n$. While the number of independent parameters for a generic unitary orbit is given by $\prod_i^n d_i - 1$. $\square$

Let me illustrate this with a simple example of a two-qubit composite system. But first, a word of caution is necessary. If an unitary orbit contains a product state that *does not* mean that all states on the that orbit are of product form. A simple example is a maximally entangled state: it is a pure state belonging to the unitary orbit of pure states, yet it is not of product form. But, of course, the unitary orbit containing the maximally entangled state will also contain pure product states.

The smallest composite system is the two-qubit system with $d = 4$. The space of the unitary orbits for the two-qubit case is confined to a tetrahedron the three dimensional space [2]. I am interested to identify the region of the tetrahedron that is occupied by unitary orbits that contain product states. To solve this problem I need to match the eigenvalues of a generic two-qubit state to the eigenvalues of a qubit state in outer product with itself

$$\text{Spec}(\rho^A) \times \text{Spec}(\rho^B) = \text{Spec}(\rho^{AB})$$

$$\begin{pmatrix} 1 + a + b + ab \\ 1 + a - b - ab \\ 1 - a + b - ab \\ 1 - a - b + ab \end{pmatrix} = \begin{pmatrix} 1 + \sqrt{6}x + \sqrt{2}y + z \\ 1 - \sqrt{6}x + \sqrt{2}y + z \\ 1 - 2\sqrt{2}y + z \\ 1 - 3z \end{pmatrix}. \quad (2)$$

Above the spectrum of a two-qubit states is given in terms of three parameters as [2, 4]. The constraint of Eq. 2 leads to equation $z^2 + z + \frac{1}{\sqrt{2}}(z-1)(\sqrt{3}x+y) + \sqrt{3}xy - y^2 = 0$ with two solutions for z . We only consider the positive solution, plotted in Fig. 2 [18]. The solution of positive z shows that the set of unitary orbits containing products states are confined to a two-dimensional plane in a three-dimensional manifold.

Note that the surface of unitary orbits containing product states runs along two of the six possible sides. The surface then divide the total space into two regions. This is a necessary condition for the plane containing unitary orbits with product states. I shall prove this now.

I would like make a few remarks about unitary orbits with product states embedded within the space all uni-




Figure 2: The geometry of unitary orbits for two qubit is represented by a tetrahedron. Above, P represents all pure states, M_2 represents states with two identical non-zero eigenvalues, M_3 represents states with three identical non-zero eigenvalues, and O represent the fully mixed state. The surface through the tetrahedron represents the set of unitary orbits that contain product states.

tary orbits. Once again, the dimensions for the unitary orbit manifold is given by the product of dimensions of each subspace, while the dimensions of the manifold that represent the unitary orbits that contain product states is given by the sum of the number of independent parameters of each subspace. The surface of unitary orbits containing product states for $d = 6$ (a qubit-qutrit system) is a 3-dimensional surface in a 5-dimensional volume. For $d = 8$ (three qubit system) case with, the surface of unitary orbits containing product states is also a 3-dimensional surface, but in a 7-dimensional volume. Roughly speaking, the set of correlated states in the three-qubit system is much higher than the correlations in the qubit-qutrit system for product states as the base measure. The space of unitary orbits grows much faster than the subspace that contains product states; this should be cautionary statement for an analysis that depends on only product states; as it may lack generality.

Next I want to look at the case of classically correlated states (not separable states), as defined by Henderson and Vedral, and independently by Ollivier and Zurek [5, 6] $\rho_{cl} = \sum_i^{i=d_A} a_i |i\rangle \langle i| \otimes \rho_i^B$. Where a_i are classical weights satisfying $\sum_i a_i = 1$ and d_A is dimensions of subsystems A .

Theorem. All bipartite-unitary orbits contain classically correlated states.

Proof. The classically correlated state is block diagonal, but it would fruitful to fully diagonalize it yielding the eigenvalues of any the unitary orbit containing it. Consider a set of unitary operators in the space of B that diagonalize the i th state, $\rho_i^B \rightarrow u_i^B \rho_i^B u_i^{B\dagger} = \sum_j b_j^{(i)} |j\rangle \langle j|$, where $b_j^{(i)}$ is the j th eigenvalues of ρ_i^B . Let me now construct the following *control-unitary* operator $U_{cd} = \sum_i |i\rangle \langle i| \otimes u_i$. Applying U to ρ_{cl} yields a diagonal

matrix.

$$\begin{aligned}
\rho_{cl,diag} &= U_{cd}\rho_{cl}U_{cd}^\dagger \\
&= \sum_{ijk} a_i |j\rangle \langle j| i\rangle \langle i| k\rangle \langle k| \otimes u_j^B \rho_i^B u_k^{B\dagger} \\
&= \sum_{ij} a_i b_j^{(i)} |ij\rangle \langle ij|. \tag{3}
\end{aligned}$$

Above $a_i b_j^{(i)}$ are the eigenvalues of state $\rho_{cl,diag}$, and therefore this is the form of the eigenvalues a classically correlated state and of the unitary orbit to which contains a classically correlated state.

Consider an arbitrary state $\rho = \sum_{ij} \lambda_{ij} |v_{ij}\rangle \langle v_{ij}|$, with eigenvalues λ_{ij} and (generally not separable) basis $|v_{ij}\rangle$, where i and j run up to the dimensions of spaces A and B respectively. By the definition of a unitary transformation there exists an operator W that connects the basis $\{|ij\rangle\}$ to the basis $\{|v_{ij}\rangle\}$, as $W|v_{ij}\rangle = |ij\rangle$. Next, rewrite the eigenvalues λ_{ij} as $\lambda_{ij} = a_i b_j^{(i)}$, where $a_i \equiv \sum_j \lambda_{ij}$ and $b_j^{(i)} \equiv \frac{\lambda_{ij}}{a_i}$. By definition both a_i and $b_j^{(i)}$ are positive numbers less than (or equal to) 1 satisfying the conditions $\sum_j b_j^{(i)} = \sum_j \frac{\lambda_{ij}}{a_i} = \frac{\sum_j \lambda_{ij}}{\sum_j \lambda_{ij}} = 1$ and $\sum_i a_i = \sum_i \sum_j \lambda_{ij} = 1$. The two conditions are the trace condition for $\rho_{cl,diag}$ and its subsystems. And finally we can connect a generic state ρ to a classically correlated state ρ_{cl} by the aid of unitary transformations U_{cd} and W .

$$\begin{aligned}
U_{cd}^\dagger W \rho W^\dagger U_{cd} &= U_{cd}^\dagger \left(\sum_{ij} a_i b_j^{(i)} |ij\rangle \langle ij| \right) U_{cd} \\
&= U_{cd}^\dagger \rho_{cl,diag} U_{cd} = \rho_{cl}. \tag{4}
\end{aligned}$$

This completes the proof that any unitary orbit containing a generic state ρ also contains a classically correlated state ρ_{cl} . $\square$

There is no real difference between the space AB and space BA apart from labeling; the result above can be extended for classical correlated states in the space of B by swapping the definitions for a_i and b_{ij} [19].

Before I conclude this paper I would like to discuss unitary orbits that contain states with entanglement briefly. Above I showed that each unitary orbit contains classically correlated states, which are unentangled. To perform this analysis on equal footing I consider the maximum entanglement an orbit can possess. Verstraete, Audenaert, and De Moor [7] have calculated the states with maximally entanglement for two qubits. They further showed that these states maximize Negativity [8], concurrence [9, 10], and relative entropy [11].

In Fig. 3 I have plotted equi-negativity surfaces within the tetrahedron. Note in the region near the fully mixed state there is no entanglement however almost all unitary orbits do not contain product states (see Fig. 2). This is a simple and method to see the visualize how entanglement is distributed in the state space. There are more




Figure 3: The three surface through the tetrahedron represent (from front to back) unitary orbits with maximum negativity of 0, $\frac{1}{4}$, and $\frac{1}{2}$.

applications of the tools developed above for studying entanglement. However I will refrain myself from discussing them presently. These topics will be discussed in greater detail in an upcoming paper [12].

The concept of unitary orbit is rather simple, yet it has allowed much insight in how multipartite systems are correlated. I showed that the set of unitary orbits containing product states is a very small subset of set of all unitary orbits. Which in return means that the set product states is a very small subset of set of all states. Any realistic model of a multipartite quantum system *should not* assume that the parts are simplify separable. That assumption leads to gross simplification which could undermine a great deal of physical effects. This was one the major goal of this paper. There has been a great deal of discussion regarding initially product state assumption in open quantum dynamics [13, 14, 15, 16]. I have not definitively shown that when initially product state assumption is retained one is restricted to a small subspace of the total state space.

The other immediate application of the calculations above is seen in the derivation of the non-Markovian master equation due Rodríguez-Rosario and Sudarshan [17]. In their derivation they start with a generic state of the system and the environment, which is assumed to be of product form at some point in its (unitary) history. This assumption cannot hold for a generic state. However, now that we know that all unitary orbits contain classically correlated states. The derivation is fully generalized by simply making the substitution of classically correlated states in place of product states in Rodríguez-Rosario and Sudarshan's derivation.

I thank Luis Boya for introducing me to this problem. I am grateful to Francesco Buscemi, Hugo Cable, Earl Campbell, Kuldeep Dixit, Elizabeth Rieper, César Rodríguez-Rosario, George Sudarshan, and Vlatko Vedral for helpful discussions. This work was financially supported by the National Research Foundation and the Ministry of Education of Singapore.

* Electronic address: cqtkkm@nus.edu.sg

- [1] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States* (Cambridge University Press, 2006).
- [2] L. Boya and K. Dixit, Phys. Rev. A **78**, 042108 (2008).
- [3] T. Tilma and E. C. G. Sudarshan, J. Phys. A **35**, 10467 (2002).
- [4] M. S. Byrd and N. Khaneja, Phys. Rev. A **68**, 062322 (2003).
- [5] L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001).
- [6] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901 (2001).
- [7] F. Verstraete, K. Audenaert, and B. De Moor, Phys. Rev. A **64**, 012316 (2001).
- [8] Peres, Phys. Rev. Lett. **77**, 1413 (1996).
- [9] S. Hill and W. K. Wootters, Phys. Rev. Lett. **78**, 5022 (1997).
- [10] W. K. Wootters, Phys. Rev. Lett. **80**, 2245 (1998).
- [11] V. Vedral and M. Plenio, Phys. Rev. A **57**, 1619 (1998).
- [12] K. Modi and V. Vedral (In preparation.).
- [13] R. Alicki, Phys. Rev. Lett. **75**, 3020 (1995).
- [14] P. Pechukas, Phys. Rev. Lett. **73**, 1060 (1994).
- [15] T. F. Jordan, A. Shaji, and E. C. G. Sudarshan, Phys. Rev. A **70**, 052110 (2004).
- [16] C. A. Rodríguez-Rosario, K. Modi, A. Kuah, E. C. G. Sudarshan, and A. Shaji, arXiv:quant-ph/0703022v3 (2007).
- [17] C. A. Rodríguez-Rosario and E. C. G. Sudarshan, arXiv:0803.1183 (2008).
- [18] I have made one additional contraction by ordering the eigenvalues from large to small (top to bottom in Eq. 2). In that sense, I have contracted twenty-four tetrahedrons (permutations of four) into one tetrahedron [2]. The negative solution therefore belongs to a space that has been contracted by permutation symmetry.
- [19] If an $a_k = 0$ for $i = k$, that means that all $\lambda_{kj} = 0$ for all j . The only requirement we have so far set is that $\lambda_{ij} = a_i b_j^{(i)}$. When the l.h.s is zero, a_i on the r.h.s is also zero, and $b_j^{(i)}$ can be anything. Another way to look at this is by considering a_i as classical wights. When the classical weight is zero then the corresponding term for the subsystem B can be anything or neglected all together.